

ON SEISMIC SIGNATURES OF RAPID VARIATION

G. Houdek¹ and D. O. Gough^{1,2}

¹*Institute of Astronomy, University of Cambridge, Cambridge CB3 0HA, UK*

²*Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge CB3 0WA, UK*

ABSTRACT

We present an improved model for an asteroseismic diagnostic contained in the frequency spacing of low-degree acoustic modes. By modelling in a realistic manner regions of rapid variation of dynamically relevant quantities, which we call acoustic glitches, we can derive signatures of the gross properties of those glitches. In particular, we are interested in measuring properties that are related to the helium ionization zones and to the rapid variation in the background state associated with the lower boundary of the convective envelope. The formula for the seismic diagnostic is tested against a sequence of theoretical models of the Sun, and is compared with seismic diagnostics published previously by Monteiro & Thompson (1998, 2005) and by Basu et al. (2004).

Key words: solar interior; solar oscillations; stellar oscillations.

1. INTRODUCTION

Abrupt variation in the stratification of a star (relative to the scale of the inverse radial wavenumber of a seismic mode of oscillation), such as that resulting from the (smooth, albeit acoustically relatively abrupt) depression in the first adiabatic exponent $\gamma_1 = (\partial \ln p / \partial \ln \rho)_s$ caused by the ionization of helium, where p , ρ and s are pressure, density and specific entropy, or from the sharp transition from radiative to convective heat transport at the base of the convection zone, induces small-amplitude oscillatory components (with respect to frequency) in the spacing of the cyclic eigenfrequencies $\nu_{n,l}$ of seismic oscillation. One might hope that the variation of the sound speed induced by helium ionization might enable one to determine from the low-degree eigenfrequencies a measure that is directly related to, perhaps even almost proportional to, the helium abundance, with little contamination from other properties of the structure of the star.

A convenient and easily evaluated measure of the oscillatory component is the second multiplet-frequency difference with respect to order n amongst modes of like degree l :

$$\Delta_2 \nu_{n,l} \equiv \nu_{n-1,l} - 2\nu_{n,l} + \nu_{n+1,l} \quad (1)$$

(Gough, 1990). Any localized region of rapid variation of either the sound speed or the density scale height, or a spatial derivative of them, which here we call an acoustic glitch, induces an oscillatory component in $\Delta_2 \nu$ with a ‘cyclic frequency’ approximately equal to twice the acoustic depth

$$\tau = \int_{r_{\text{glitch}}}^R c^{-1} dr \quad (2)$$

of the glitch, and with an amplitude which depends on the amplitude of the glitch and which decays with ν once the inverse radial wavenumber of the mode becomes comparable with or greater than the radial extent of the glitch.

We report here on a model formula for the oscillatory contribution to the second frequency difference that is directly related to the gross properties of the glitch, and compare how well it fits actual eigenfrequency differences with corresponding fits of previously published formulae.

2. THE FITTING FUNCTION

For a nonrotating star with vanishing pressure gradient at the surface, the angular eigenfrequencies ω of adiabatic oscillation obey a variational principle (e.g. Chandrasekhar 1963) which can be used to estimate the contribution $\delta\omega$ to ω from the acoustic glitch. In estimating the contribution from helium ionization we retain only the dominant glitch term ($\delta\gamma_1/\gamma_1$ at fixed pressure and temperature), and we represent $\delta\gamma_1/\gamma_1$ by a pair of (negative) Gaussian functions of τ with widths Δ_I and Δ_{II} , whose integrals are Γ_I and Γ_{II} , and which are centred about the acoustic depths τ_I and τ_{II} of the first and second ionization zones of helium beneath the seismic surface $r = R$ of the star. After converting to cyclic frequency $\nu = \omega/2\pi$, the oscillatory component is given asymptotically by

$$\begin{aligned}\delta_{\gamma,\text{osc}}\nu &\simeq -\Gamma_{\text{II}}\nu_0 \left[\nu + \frac{1}{2}(m+1)\nu_0 \right] \\ &\times (\mu\beta\kappa_{\text{I}}^{-1}e^{-8\pi^2\mu^2\kappa_{\text{I}}^2\Delta_{\text{II}}^2\nu^2}\cos 2\psi_{\text{I}} \\ &+ \kappa_{\text{II}}^{-1}e^{-8\pi^2\kappa_{\text{II}}^2\Delta_{\text{II}}^2\nu^2}\cos 2\psi_{\text{II}}) \quad (3)\end{aligned}$$

(Houdek & Gough, 2007). We evaluate the phases $\psi_{\text{I}} = \psi(\tilde{\tau}_{\text{I}})$ and $\psi_{\text{II}} = \psi(\tilde{\tau}_{\text{II}})$, where $\omega\tilde{\tau} = \omega\tau + \epsilon_{\text{II}}$, by representing the envelope by a plane-parallel polytrope of index $m = 3.5$ and adding a phase constant ϵ_{II} to $\omega\tau$ to account for the deviation of the actual envelope from the polytrope:

$$\psi(\tau) = \omega\tau\kappa - (m+1)\cos^{-1}\left(\frac{m+1}{\tau\omega}\right) + \frac{\pi}{4}. \quad (4)$$

In equation (3), $\kappa_{\text{I}} = \kappa(\tilde{\tau}_{\text{I}})$ etc, with $\kappa(\tau) = [1 - (m+1)^2/4\pi^2\nu^2\tau^2]^{1/2}$; also ν_0 is a characteristic value of the first frequency difference $\Delta_{\text{I}}\nu$ whose value is half the inverse total acoustic radius T of the star: $\nu_0 = \omega_0/2\pi = 1/2T$. We have found that the ratios $\beta = \Gamma_{\text{I}}\Delta_{\text{II}}/\Gamma_{\text{II}}\Delta_{\text{I}}$, $\eta = \tau_{\text{I}}/\tau_{\text{II}}$ and $\mu = \Delta_{\text{I}}/\Delta_{\text{II}}$ hardly vary amongst stellar models whose masses and radii vary by factors of at least five. We have accordingly adopted constant values, namely $\beta = 0.45$, $\eta = 0.7$ and $\mu = 0.9$ for solar-like stars.

To obtain a complete description of $\Delta_2\nu$ we must add an oscillatory contribution from the near discontinuity in the density scale height at the base τ_c of the convection zone:

$$\begin{aligned}\delta_{\text{c,osc}}\nu &\simeq \hat{A}_c\nu_0^3\nu^{-2} (1 + 1/16\pi^2\tau_0^2\nu^2)^{-1/2} \\ &\times \{\cos[2\psi_c + \tan^{-1}(4\pi\tau_0\nu)]\}, \quad (5)\end{aligned}$$

with

$$\hat{A}_c = \frac{c_c^2}{8\pi\omega_0^2} [d^2 \ln \rho / dr^2]_{\tau_c}^{r_c+}, \quad (6)$$

where $\psi_c = \psi(\tau_c + \omega^{-1}\epsilon_c)$ is the polytropic phase at the base of the convection zone, the phase constant ϵ_c being allowed to differ from ϵ_{II} to account for the fact that the region of the convection zone beneath the helium ionization zones is not an exact polytrope; the quantity τ_0 is a characteristic scale over which the sound speed in the radiative interior relaxes to a smooth extrapolation from the convection zone (and which we also take to be constant, 80 s, for all the stars we consider).

From these expressions it is straightforward to evaluate the second difference $\Delta_2\nu$, to which must be added a smooth term which we represent by a third-degree polynomial in ν^{-1} :

$$\Delta_{2,\text{sm}} = \sum_{i=0}^3 a_i \nu^{-i}. \quad (7)$$

The eleven parameters A_{II} , A_c , Δ_{II} , τ_{II} , τ_c , ϵ_{II} , ϵ_c and a_i are adjusted to fit by least squares the theoretical curve to

the second frequency differences of the actual eigenfrequencies of the modes.

We compare the current seismic diagnostic (3)–(7) with diagnostics published previously by Monteiro & Thompson (1998, 2005) and by Basu et al. (2004).

2.1. The seismic diagnostic by Monteiro & Thompson

This diagnostic is based on a single-triangular approximation to $\delta\gamma_1/\gamma_1$ due to He ionization. The resulting formula for the frequency perturbation is

$$\begin{aligned}\delta_{\text{osc}}\nu &\simeq b_1 \frac{\sin^2(2\pi\nu\hat{\beta})}{\nu\hat{\beta}} \cos(4\pi\nu\tau_{\text{II}} + 2\epsilon_{\text{II}}) \\ &+ \left[\frac{c_1^2}{\nu^4} + \frac{c_2^2}{\nu^2} \right]^{1/2} \cos(4\pi\nu\tau_c + 2\epsilon_c), \quad (8)\end{aligned}$$

in which $\hat{\beta}$ is the half-width and b_1 is proportional to the height of the triangle, from which second differences can be computed. We set to zero the fitting parameter c_2 , a nonzero value of which, in the original formulation, would account for an putative adiabatic extension to the convection zone produced by overshooting (which is not present in our solar models). By so doing, the seismic diagnostic (8) has the same number, 11, of fitting parameters as the seismic diagnostic (3)–(7) when adopting the third-degree polynomial (7) to represent the smooth contribution.

2.2. The seismic diagnostic by Basu et al.

In this formulation the seismic diagnostic is that used by Basu (1997), which was invented by Basu, Antia and Narasimha (1994) for studying the base of the convection zone. Ballot et al. (2004) and Piau et al. (2005) have adopted essentially the same formula. It is not directly linked to the form of the ionization-induced acoustic glitch, but instead was motivated by the oscillatory signature of a discontinuity in sound speed, whose amplitude was modified to make it decay with frequency approximately in the manner of the second differences of the eigenfrequencies. The functional form is given by

$$\begin{aligned}\Delta_2\delta_{\text{osc}}\nu &\simeq \left(b_1 + \frac{b_2}{\nu^2} \right) \cos(4\pi\nu\tau_{\text{II}} + 2\epsilon_{\text{II}}) \\ &+ \left(c_1 + \frac{c_2}{\nu^2} \right) \cos(4\pi\nu\tau_c + 2\epsilon_c). \quad (9)\end{aligned}$$

Here we set $c_1 = 0$, to remove the effect of the erroneous sound-speed discontinuity at the base of the convection zone and to render the functional form of the contribution at $\tau = \tau_c$ (nearly) consistent with those in the other two signatures (5) and (8); and we adopt the third-degree polynomial (7) to represent the smooth contribution, thereby using the same number (11) of fitting parameters as the seismic diagnostics (3)–(7) and (8), (7).




Figure 1. Denotation of the nine calibrated solar models which are used for testing the formulation of the second frequency differences. The ‘central model’ is model 0; the set of the four models (1)–(4) have a constant value of Z but varying age; the second set of the models, (5)–(8), have constant age but varying Z .

3. TEST MODELS

Two sets of calibrated evolutionary models of the Sun, computed by Gough and Novotny (1990), were used. One set of models has a constant value for the heavy-element abundance Z but with age varying in uniform steps of approximately 0.051 in $\ln t$, and the other set has constant age but varying heavy-element abundance in uniform steps of approximately 0.016 in $\ln Z$ (see Fig. 1).

4. RESULTS

All three diagnostics were fitted by the same numerical algorithm. The lower panels of Figs 2–4 show the results for the three seismic diagnostics (3)–(7), (8) and (7), and (9) and (7), fitted by least-squares to the adiabatically computed eigenfrequencies of the nine test models. Plotted are the frequency-averaged amplitudes, $\langle A_{II} \rangle$ and $\langle A_c \rangle$, of the two oscillatory seismic contributions associated with He II ionization and with the base of the convection zone, obtained by averaging the appropriate calibrated formula uniformly over the frequency range of the data plotted in the upper panels. Also plotted are the values for the acoustic depths τ_{II} and τ_c . Depending on the seismic diagnostic, the values of either Δ_{II} , $\hat{\beta}$ or the phase constant ϵ_{II} are shown, and also the standard measure χ^2 of the goodness of the fit between $\Delta_2\nu$ of the modelled frequencies and the calibrated seismic diagnostic. In all three cases the frequency-averaged amplitude $\langle A_{II} \rangle$ increases with the helium abundance Y of the test models; moreover, the amplitudes Γ_{II} and b_1 inferred from diagnostics (3)–(7) and (8), (7) respectively also increase with Y (Γ_{II} varying almost linearly in the Z -varying sequence, b_1 varying almost linearly in the t -varying sequence), the latter result being apparently

at variance with the findings of Monteiro and Thompson (2005). The seismic diagnostic (3)–(7) fits $\Delta_2\nu$ of the model eigenfrequencies the best, as is evinced by its smallest values of χ^2 .

The quality of the fits is also illustrated in the upper panels of Figs 2–4, which show the second frequency differences over the whole frequency range considered for the central model 0. It is also clear from comparing these figures that the seismic diagnostic (3)–(7) reproduces $\Delta_2\nu$ of the model frequencies the most faithfully.

It is interesting to note that in determining the parameters in the formulae the fitting procedure gives greater weight to the low-frequency end of the ionization-induced component $\Delta_{2,\gamma}$, where the amplitude is the greatest, and greater weight to the high-frequency end of the contribution $\Delta_{2,c}$ from the base of the convection zone, where misfits cannot be offset by adjustments to $\Delta_{2,\gamma}$; therefore, the diagnostics (8), (7) and (9), (7) fail principally in the mid-range. Some large-scale discrepancy can be seen also at high frequency for formula (9), (7) because $\Delta_{2,\gamma}$ decays with frequency too slowly. We acknowledge that these discrepancies might be either smaller or hardly greater than the random errors in imminent asteroseismic data; nevertheless, fitting an erroneous formula can lead to the inferences from that fitting being biased, and should therefore be avoided.

ACKNOWLEDGEMENTS

GH is grateful for support from the Particle Physics and Astronomy Research Council.

REFERENCES

- [1] Ballot J., Turck-Chièze, S., García R.A., 2004, A&A, 423, 1051
- [2] Basu S., 1997, MNRAS, 288, 527
- [3] Basu S., Antia H., Narasimha D., 1994, MNRAS, 267, 209
- [4] Basu S., Mazumdar A., Antia H.M., Demarque P., 2004, MNRAS 350, 277
- [5] Chandrasekhar S., 1963, ApJ 138, 896
- [6] Gough D.O., 1990, in Progress of Seismology of the Sun and Stars, Lecture Notes in Physics, Vol. 367, Y. Osaki, H. Shibahashi, eds. Springer Verlag, p. 283
- [7] Gough D.O. Novotny, E., 1990, Solar Phys., 128, 143
- [8] Houdek G., Gough D.O. 2007, MNRAS, in press
- [9] Monteiro M.J.P.F.G., Thompson M., 1998, in Deubner F.-L., Christensen-Dalsgaard J., Kurtz D., eds. Proc. IAU Symp.185, New Eyes to See Inside the Sun and Stars. Kluwer, Dordrecht, p. 317
- [10] Monteiro M.J.P.F.G., Thompson M., 2005, MNRAS, 361, 1187
- [11] Piau L. Ballot J. Turck-Chièze S., 2005, A&A, 430, 571




Figure 2. Top: The symbols are second differences $\Delta_2\nu$, defined by equation (1), of low-degree frequencies obtained from adiabatic pulsation calculations of the central model 0. The curve is the seismic diagnostic (3)–(7) whose eleven parameters have been adjusted to fit the data by least squares. Bottom: Model properties determined from the seismic diagnostic (3)–(7) for the nine test models. Models with varying age are connected by solid lines, those with varying Z by dashed lines. Some model identifiers are written in the upper left panel. The smallest (central) frequency value used in the least-square fitting is $\nu \simeq 1322 \mu\text{Hz}$, the largest is $\nu \simeq 4058 \mu\text{Hz}$. The lower right panel is the standard measure χ^2 for the goodness of the least-squares fit.




Figure 3. Top: The symbols are second differences $\Delta_2\nu$, defined by equation (1), of low-degree frequencies obtained from adiabatic pulsation calculations of the central model 0. The curve is the seismic diagnostic (8),(7) whose eleven parameters have been adjusted to fit the data by least squares. Bottom: Model properties determined from the seismic diagnostic (8),(7) for the nine test models. The smallest (central) frequency value used in the least-square fitting is $\nu \simeq 1322 \mu\text{Hz}$, the largest is $\nu \simeq 4058 \mu\text{Hz}$. The lower right panel is the standard measure χ^2 for the goodness of the least-squares fit.




Figure 4. Top: The symbols are second differences $\Delta_2\nu$, defined by equation (1), of low-degree frequencies obtained from adiabatic pulsation calculations of the central model 0. The curve is the seismic diagnostic (9),(7) whose eleven parameters have been adjusted to fit the data by least squares. Bottom: Model properties determined from the seismic diagnostic (9),(7) for the nine test models. The smallest (central) frequency value used in the least-square fitting is $\nu \simeq 1322 \mu\text{Hz}$, the largest is $\nu \simeq 4058 \mu\text{Hz}$. The lower right panel is the standard measure χ^2 for the goodness of the least-squares fit.